## Pulmonary and Sleep Associates of Marin



100 Rowland Way, Suite 300 Novato, CA 94945

Phone: 415-878-0225 Fax: 415-878-0215

MarinPulmonarySleep.com

DATE: NAME:

## Assess the Severity of Restless Legs Syndrome

## USING THE INTERNATIONAL RLS (IRLS) RATING SCALE

The IRLS Rating Scale was developed by the International Restless Legs Syndrome Study Group (IRLSSG) to assess the severity of a patient's RLS symptoms. Ask your patient the 10 questions listed below and have them rate their symptoms from 0 to 4. Mark each answer and total their IRLS score. It will range from 0 to 40.

The IRLS Rating Scale was validated in a controlled study and found to have high levels of internal consistency, interexaminer reliability, and test-retest reliability. The IRLSSG is an organization of professionals committed to advancing basic and clinical research on RLS. To learn more about the group, visit www.irlssg.org.

In the past week:	Score
Overall, how would you rate the RLS discomfort in your legs or arms?  4 Very severe 3 Severe 2 Moderate 1 Mild 0 None	
Overall, how would you rate the need to move around because of your RLS symptoms?  4 Very severe 3 Severe 2 Moderate 1 Mild 0 None	
Overall, how much relief of your RLS arm or leg discomfort did you get from moving around?  4 No relief 3 Mild relief 2 Moderate relief 1 Either complete or almost complete relief 0 No RLS symptoms to be relieved	
How severe was your sleep disturbance from your RLS symptoms? 4 Very severe 3 Severe 2 Moderate 1 Mild 0 None	
How severe was your tiredness or sleepiness during the day due to your RLS symptoms? 4 Very severe 3 Severe 2 Moderate 1 Mild 0 None	
How severe was your RLS as a whole? 4 Very severe 3 Severe 2 Moderate 1 Mild 0 None	
How often did you get RLS symptoms? 4 Very often (6–7 days in 1 week). 3 Often (4–5 days in 1 week). 2 Sometimes (2–3 days in 1 week). 1 Occasionally (1 day in 1 week). 0 Never	
When you had RLS symptoms, how severe were they on average? 4 Very severe (8 h or more per 24 h) 3 Severe (3–8 h per 24 h) 2 Moderate (1–3 h per 24 h) 1 Mild (less than 1 h per 24 h) 0 None	
Overall, how severe was the impact of your RLS symptoms, on your ability to carry out your daily affairs, for example, carrying out a satisfactory family, home, social, school, or work life?  4 Very severe 3 Severe 2 Moderate 1 Mild 0 None	
How severe was your mood disturbance from your RLS symptoms—for example angry, depressed, sad, anxious, or irritable?  4 Very severe 3 Severe 2 Moderate 1 Mild 0 None	
Add the scores from above and share the total with your patient.  Mild 0 to 10 Moderate 11 to 20 Severe 21 to 30 Very Severe 31 to 40	Total Score